

CRISPR/CAS9-MEDIATED KNOCK-OUT OF TOMATO DMR6 GENE TO CONFER TOLERANCE AGAINST PATHOGENS

MAIOLI A.*, MILORO F.***, GIANOGLIO S.***, LI R.*, VALENTINO D.*, MILANI A. M.*, COMINO C.*, LANTERI S.*, ACQUADRO A.*, MOGLIA A.*

*) Department of Agricultural, Forest and Food Science, University of Turin, Italy

**) Department of Plant Biotechnology, Agricultural Biotechnology Institute, National Agricultural Research and Innovation Centre, 2100 Gödöllő, Hungary

***) Instituto De Biología Molecular y Celular De Plantas (IBMCP) CSIC-UPV, Ingeniero Fausto Elio/n, 46022 Valencia, Spain

DMR6, susceptibility, CRISPR/Cas9, pathology, tomato

A promising strategy to confer broad-spectrum and long-lasting resistance against pathogens in plants is based on the disabling of susceptibility genes (S-genes). These genes, required by the pathogen to infect a plant at some step of the infection process, support plant/pathogen compatibility. The S-gene Downy Mildew Resistance 6 (DMR6), which is related to salicylic acid catabolism, negatively regulates defence genes' expression. In Arabidopsis and potato, DMR6 confers tolerance against *Phytophthora capsici* and *Phytophthora infestans*, thus we assessed its role of DMR6 in affecting in tomato the susceptibility against late blight (LB) in tomato caused by *P. infestans*.

Two homologues (dmr6-1 and dmr6-2) were identified in tomato and, based on their transcription activation in response to *P. infestans*, dmr6-1 was selected as the most promising target. In order to induce knock-out mutations, a multiplexing CRISPR/Cas9 approach based on three gRNAs designed on dmr6-1 was applied.

The experimental work led to the production of five in vitro plantlets. Illumina deep sequencing of the target sites revealed the successful editing of dmr6 in all of them, with the insertion of 1bp as the most frequent observed mutation. After selfing, mutated alleles were stably inherited in ten T1 progeny. One plant was found carrying mutations in homozygous state in all the target regions and, due to the segregation of Cas9, the absence of transgene. A whole genome re-sequencing (Illumina) of this mutant was also performed to verify the absence of off-target genomic effects.

T1 mutants were then evaluated through a Detached-Leaf Assay (DLA) using *P. infestans* and all of them showed reduced disease symptoms if compared with their wild types. The tolerance of mutant lines against *O. neolycopersici*, *B. cinerea*, *A. alternata* and *C. fulvum* will be tested in future experiments.