Proceedings of the LXVI SIGA Annual Congress Bari, 5/8 September, 2023 ISBN: **978-88-944843-4-2** 

Oral Communication Abstract – 1.05

## DEVELOPMENT AND APPLICATION OF SINGLE PRIMER ENRICHMENT TECHNOLOGY (SPET) SNP ASSAY FOR POPULATION GENOMICS ANALYSIS AND CANDIDATE GENE DISCOVERY IN LETTUCE

TRIPODI P.\*, BERETTA M.\*\*, PELTIER D.\*\*\*, KALFAS I.\*\*\*\*, VASILIKIOTIS C.\*\*\*\* , LAIDET A.\*\*\*\*, BRIAND G.\*\*\*\*, AICHHOLZ C.\*\*\*\*\*, ZOLLINGER T.\*\*\*\*\*\*, VAN TREUREN R.\*\*\*\*\*\*, SCAGLIONE D.\*\*\*\*\*\*\*, GORITSCHNIG S.\*\*\*\*\*\*\*\*

\*) CREA Research Centre for Vegetable and Ornamental Crops, Via dei Cavalleggeri 25, 84098, Pontecagnano Faiano, SA, Italy \*\*\*) ISI Sementi SpA, 43036 Fidenza (PR), Italy \*\*\*) LIMAGRAIN - Vilmorin-Mikado, Route du Manoir, 49250 La Ménitré, France \*\*\*\*\*) American Farm School, Thessaloniki, Greece \*\*\*\*\*) Gautier Semences Route d'Avignon 13630 Eyragues, France \*\*\*\*\*\*) Sativa Rheinau AG, Switzerland \*\*\*\*\*\*\*) Zollinger Conseilles Sarl, Switzerland \*\*\*\*\*\*\*) Centre for Genetic Resources, the Netherlands (CGN), Wageningen University and Research, P.O. Box 16, 6700 AA Wageningen, the Netherlands \*\*\*\*\*\*\*) IGA Technology Services Srl, Udine, Italy \*\*\*\*\*\*\*\*) ECPGR Secretariat c/o Alliance of Bioversity International and CIAT, Via dei Tre Denari 472A 00054 Maccarese (RM)Via di San Domenico 1, 00153 Rome, Italy

lettuce, high-throughput genotyping, genomic diversity, GWAS, candidate
genes

Cultivated lettuce (*Lactuca sativa* L.) is a main leafy vegetable crop widely appreciated by consumers for the content of fibres and the lowcalorie intake being furthermore a rich source of vitamin C, iron, folate as well as different health-beneficial bioactive compounds. Recent years witnessed astonishing advancements in the development of cutting-edge technologies for next-generation sequencing opening new frontiers for dissecting the genetic basis of crops and discovery novel genes of interest. Among these, SPET (single primer enrichment technology) is a novel high-throughput genotyping method based on short-read sequencing of specific genomic regions harboring polymorphisms. SPET provides an

efficient and reproducible method for genotyping target loci, overcoming the limits associated with other reduced representation library sequencing random methods that are based on а sampling of genomic loci. The possibility to sequence regions surrounding a target SNP allows the discovery of thousands of closely linked, novel SNPs. Here we report the design and application of the first SPET panel in lettuce, consisting of 41,547 probes spanning the whole genome and designed to target both coding (~96%) and intergenic (~4%) regions. A total of 81,531 SNPs were surveyed in 160 lettuce accessions originating from a total of 10 countries in Europe, America and Asia and representing ten different horticultural types. Model ancestry population structure clearly separated the cultivated accessions (Lactuca sativa) from accessions of its presumed wild progenitor (L. serriola), revealing a total of six genetic subgroups, that reflected a differentiation based on cultivar typology. Phylogenetic relationships and principal component analysis revealed a clustering of butterhead types and a general differentiation between germplasm originating from Western and Eastern Europe. To determine the potentiality of SPET for gene discovery, we performed genome wide association analysis for main agricultural traits in *L. sativa* using eight models (GLM naive, GLM-Q, GLM-PCA, MLM, MLMM, compare their CMLM, FarmCPU and BLINK) to strength and power for association detection. Robust associations were detected for seeds colour on chromosome 7 at 50 Mbp. Colocalization of association signals was found for outer leaf colour and leaf anthocyanin content on the bottom chromosome 9 at 152 Mbp and on chromosome 5 at 96 Mbp. The association for bolting time was detected with the BLINK, farm CPU and MLM model on chromosome 7 at Detected associations fall in chromosomal regions previously 164 Mbp. hold candidate genes of interest for these traits, reported to thus confirming the effectiveness for GWAS. Our research demonstrates the reliability of SPET for detecting thousands of variable sites that help to dissect the genetic diversity of germplasm collections and to identify candidate genes for traits of agricultural interest, enabling a better use of lettuce germplasm.