Proceedings of the LXVI SIGA Annual Congress Bari, 5/8 September, 2023 ISBN: **978-88-944843-4-2**

Poster Communication Abstract - 1.20

INVESTIGATION ON THE GENETIC DIVERSITY OF GRAPEVINE CLONES FROM APULIAN AND CAMPANIAN VARIETIES

PROCINO S.***, VILLANO C.***, CARPUTO D.***, D'AGOSTINO N.****, DI SERIO E.***, FANELLI V.*, LA NOTTE P.****, MIAZZI M. M.*, MONTEMURRO C.********, TARANTO F.**, AVERSANO R.***

*) Department of Soil, Plant, and Food Sciences, Faculty of Agricultural Science, University of Bari Aldo Moro, Italy **) CNR-IBBR Institute of Bioscience and BioResources, Italy ***) Department of Agricultural Sciences, University of Naples Federico II, Italy ****) Institute for the Sustainable Protection of Plants, Bari branch, Institute for Sustainable Plant Protection, National Research Council (CNR), Italy *****) Spin-Off Sinagri, Università degli Studi di Bari Aldo Moro, Italy

Vitis vinifera clones, GBS, ddRAD, SNP markers, divergent loci

The genetic diversity of the cultivated grapevine (*Vitis vinifera* L.) arises from a large population of varieties, with sexual reproduction being contributor to inter-varietal variability. major Additionally, а spontaneous somatic mutations have played a role in shaping the vast array grape varieties. Over the centuries, farmers have collected of and propagated various selections of grapevine, labelling them as "clones". These clones often represent underexplored genetic resources that can be used to achieve successful conservation strategies and improve economically important traits. This study aimed to investigate the population structure and detect divergent loci in the genomes of 138 clones of six grapevine varieties from Campania (AGLIANICO LASCO, CAMAIOLA, and GRECO BIANCO) and Apulia (MALVASIA NERA, MINUTOLO and UVA DI TROIA). The samples from Campania and Apulia were genotyped using double-digested restriction siteassociate DNA sequencing (ddRADseq) and Genotyping-by-sequencing (GBS), respectively. In both cases, the Pinot Noir PN40024 reference genome was for calling single nucleotide polymorphisms (SNPs). used Population genetic analysis indicated that all the varieties grouped their own clones,

except for AGLIANICO LASCO which was separated into two sub-populations (AL1 and AL2). AL2 clones can be considered the true AGLIANICO LASCO, whereas AL1 individuals are homonyms of AGLIANICO. Genetic differentiation among the seven populations was further investigated by fixation index (*Fst*). At *Fst* \geq 0.80, a total of 1,016 divergent SNPs were detected, of which 285 were non redundant. Among the latter, 200 were in gene regions. The largest number of divergent SNPs was found in CAMAIOLA (N. = 453), while AL1 showed the lowest number (N. = 117). CAMAIOLA also had the largest number of divergent states than the other varieties (> 81%).

In this study, we have shown that merging different SNP datasets is possible and valuable to study the inter- and intra-specific genetic diversity of grapevine populations. This provided a repeatable framework to streamline future computational studies based on retrieving information from partial analyses performed at different times and with different techniques.

Our results also demonstrated the value of advanced genomic methods in the study of synonymy/homonymy detection (as exemplified by AGLIANICO LASCO). Finally, signatures of divergence were identified among cultivars and between Campanian and Apulian accessions, some of which were involved in environmental adaptation and linked to key traits useful for grapevine breeding, such as budding, flowering and berry ripening time. These findings emphasize the importance of exploiting the vast untapped genetic variability that characterizes the minor grape varieties and the urgent need to conserve their genetic resources using them more effectively in breeding programs.