Proceedings of the LXIV SIGA Annual Congress Online, 14/16 September, 2021 ISBN: **978-88-944843-2-8** 

Poster Communication Abstract - 7.12

## CRISPR/CAS9-MEDIATED MUTAGENESIS TO IMPROVE THE NUTRITIONAL QUALITY OF TOMATO FRUITS.

OLIVIERI F.\*, FRANCESCA S.\*, SVEVI F.\*\*, CHIAIESE P.\*, BARONE A.\*, D'AGOSTINO N.\*, RIGANO M. M.\*

 \*) Dipartimento di Agraria, Università degli Studi di Napoli Federico II, Portici
\*\*) Agenzia nazionale per le nuove tecnologie, l'energia e lo sviluppo economico sostenibile (ENEA), Centro Ricerche Casaccia, Roma

ascorbic acid, gene knock-out, tomato nutritional quality, new plant breeding techniques

Tomato (Solanum lycopersicum L.) berries are a rich source of antioxidants and other bioactive compounds, including Ascorbic Acid (AsA), which plays a key role in human health. Plants produce AsA through several biochemical pathways and many genes and enzymes in those pathways have been identified in tomato. About 25 years ago introgression lines (ILs) or sub-lines were produced, in which genomic regions of S. lycopersicum cv. M82 were replaced with the corresponding segments of the wild S. pennellii genome. In our laboratory, several sub-lines have been produced and characterized that accumulate high content of AsA in the fruits. Recently, we identified a gene in the wild region of the IL7-3 that encodes an ascorbate oxidase (AO) and that regulates the accumulation of AsA in tomato fruits. Further investigations on ILs have allowed demonstrating that pectin methylesterases (PMEs) and pectin methylesterases inhibitors (PMEIs) play a fundamental role in the biosynthesis of AsA using the D-galacturonic acid derived from pectins as metabolic precursor.

Within this motivating context, we would like to exploit the great potential of the genome editing techniques to accelerate breeding programs with the aim of improving the organoleptic and nutritional (increase in AsA content) quality of tomato fruits. To achieve our goals, we selected three target genes, potentially involved in the AsA accumulation and encoding an ascorbate oxidase, a laccase (LAC) and a PMEI, respectively. We designed

single-guide RNA molecules for all candidate genes and constructed CRISPR/Cas9 plasmids knock-out experiments. for We used different strategies to design sgRNA expression plasmids: LAC and AO were targeted with four individual sgRNAs, while PMEI with a single guide. Genetic transformation of tomato cotyledons (Red Setter and MonevMaker) was mediated by A. tumefaciens strain LBA4404. At present, we obtained two Cas9+ plants for the LAC gene and seven for the AO gene. The Cas9+ plants were micropropagated and *in vivo* transferred to obtain T1 seeds. Sanger 0ther performed to confirm gene knock-out. sequencing was genetic transformations are ongoing for these two target genes in order to increase the number of successfully edited plants. As for PMEI, we obtained 34 regenerated explants on selection media and the validation of the Cas9+ events is ongoing. In the near future, morpho-physiological and biochemical analysis (i.e., plant phenotyping; evaluation of the AsA content and total antioxidant activity) on the fruits harvested from the mutated plants will be performed to verify the improvement in the nutritional quality of tomato fruits.